

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

Keflex 125 mg/5 ml Powder for Oral Suspension

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each bottle when prepared as directed, contains as the active ingredient cefalexin equivalent to 125 mg / 5 ml cefalexin anhydrous.

Excipient: Contains 3.119g of sucrose per 5ml.

Contains allura red AC (E129).

For a full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Powder for oral suspension

White, free flowing powder.

4 CLINICAL PARTICULARS

4.1 Therapeutic Indications

Cefalexin is a semisynthetic cephalosporin antibiotic for oral suspension.

Cefalexin is indicated in the treatment of the following infections due to susceptible micro-organisms:

Respiratory tract infections.

Otitis media.

Skin and soft tissue infections.

Bone and joint infections.

Genito-urinary tract infections, including acute prostatitis.

Dental infections.

4.2 Posology and method of administration

Keflex is administered orally.

Adults: The adult dosage ranges from 1-4g daily, in divided doses; most infections will respond to a dosage of 500mg every 8 hours. For skin and soft tissue infections, streptococcal pharyngitis and mild, uncomplicated urinary tract infections, the usual dosage is 250mg every 6 hours, or 500mg every 12 hours.

For more severe infections or those caused by less susceptible organisms, larger doses may be needed. If daily doses of Keflex greater than 4g are required, parenteral cephalosporins, in appropriate doses, should be considered.

The elderly and patients with impaired renal function: As for adults. Reduce dosage if renal function is markedly impaired (see 'Special warnings').

Children: The usual recommended daily dosage for children is 25-50mg/kg (10-20mg/lb), in divided doses. For skin

and soft tissue infections, streptococcal pharyngitis and mild, uncomplicated urinary tract infections, the total daily dose may be divided and administered every 12 hours. For most infections the following schedule is suggested:

<i>Children under 5 years:</i>	125mg every 8 hours.
<i>Children 5 years and over:</i>	250mg every 8 hours.

In severe infections, the dosage may be doubled. In the therapy of otitis media, clinical studies have shown that a dosage of 75 to 100mg/kg/day, in 4 divided doses, is required.

In the treatment of beta-haemolytic streptococcal infections, a therapeutic dose should be administered for at least 10 days.

4.3 Contraindications

Cefalexin is contra-indicated in patients with known allergy to the cephalosporin group of antibiotics.

4.4 Special warnings and precautions for use

Before instituting therapy with cefalexin, every effort should be made to determine whether the patient has had previous hypersensitivity reactions to the cephalosporins, penicillins or other drugs. Cefalexin should be given cautiously to penicillin-sensitive patients. There is some clinical and laboratory evidence of partial cross-allergenicity of the penicillins and cephalosporins. Patients have had severe reactions (including anaphylaxis) to both drugs.

Pseudomembranous colitis has been reported with virtually all broad-spectrum antibiotics, including macrolides, semisynthetic penicillins and cephalosporins. It is important, therefore, to consider its diagnosis in patients who develop diarrhoea in association with the use of antibiotics. Such colitis may range in severity from mild to life-threatening. Mild cases of pseudomembranous colitis usually respond to drug discontinuance alone. In moderate to severe cases, appropriate measures should be taken.

If an allergic reaction to cefalexin occurs, the drug should be discontinued and the patient treated with the appropriate agents.

Prolonged use of cefalexin may result in the overgrowth of non-susceptible organisms. Careful observation of the patient is essential. If superinfection occurs during therapy, appropriate measures should be taken.

Cefalexin should be administered with caution in the presence of markedly impaired renal function. Careful clinical and laboratory studies, including monitoring the serum levels of cefalexin, should be made because safe dosage may be lower than that usually recommended. If dialysis is required for renal failure, the daily dose of cefalexin should not exceed 500mg.

Concurrent administration with certain other drug substances, such as aminoglycosides, other cephalosporins, or furosemide (frusemide) and similar potent diuretics, may increase the risk of nephrotoxicity.

Positive direct Coombs' tests have been reported during treatment with the cephalosporin antibiotics. In haematological studies, or in transfusion cross-matching procedures when antiglobulin tests are performed on the minor side, or in Coombs' testing of newborns whose mothers have received cephalosporin antibiotics before parturition, it should be recognised that a positive Coombs' test may be due to the drug.

A false positive reaction for glucose in the urine may occur with Benedict's or Fehling's solutions or with copper sulphate test tablets. Cefalexin may interfere with the alkaline picrate assay for creatinine.

Patients with rare hereditary problems of fructose intolerance, glucose-galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine.

4.5 Interaction with other medicinal products and other forms of interaction

As with other beta-lactam drugs, renal excretion of cefalexin is inhibited by probenecid.

In healthy subjects given single 500mg doses of cefalexin and metformin, plasma metformin C_{\max} and AUC increased by an average of 34% and 24%, respectively, and metformin renal clearance decreased by an average of 14%. No information is available about the interaction of cefalexin and metformin following multiple dose administration.

Hypokalaemia has been described in patient taking cytotoxic drugs for leukaemia when they were given gentamicin and cephalixin.

4.6 Fertility, pregnancy and lactation

Usage in pregnancy: Although laboratory and clinical studies have shown no evidence of teratogenicity, caution should be exercised when prescribing for the pregnant patient.

Usage in nursing mothers: The excretion of cefalexin in human breast milk increased up to 4 hours following a 500mg dose. The drug reached a maximum level of 4 micrograms/ml, then decreased gradually and had disappeared 8 hours after administration. Caution should be exercised when cefalexin is administered to a nursing woman, since the neonate is presented with the risk of candidiasis and CNS toxicity due to immaturity of the blood–brain barrier. There is a theoretical possibility of later sensitisation.

4.7 Effects on ability to drive and use machines

Keflex has no known influence on the ability to drive and use machines.

4.8 Undesirable effects

Gastro–intestinal: Symptoms of pseudomembranous colitis may appear either during or after antibiotic treatment. Nausea and vomiting have been reported rarely. The most frequent side–effect has been diarrhoea. It was very rarely severe enough to warrant cessation of therapy. Dyspepsia and abdominal pain have also occurred. As with some penicillins and some other cephalosporins, transient hepatitis and cholestatic jaundice have been reported rarely.

Hypersensitivity: Allergic reactions have been observed in the form of rash, urticaria, angioedema and, rarely, erythema multiforme, Stevens–Johnson syndrome and toxic epidermal necrolysis. These reactions usually subsided upon discontinuation of the drug, although in some cases supportive therapy may be necessary. Anaphylaxis has also been reported.

Haemic and Lymphatic System: Eosinophilia, neutropenia, thrombocytopenia, and haemolytic anaemia have been reported.

Other: These have included genital and anal pruritus, genital candidiasis, vaginitis and vaginal discharge, dizziness, fatigue, headache, agitation, confusion, hallucinations, arthralgia, arthritis and joint disorder. Reversible interstitial nephritis has been reported rarely. Slight elevations in AST and ALT have been reported.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via:

HPRA Pharmacovigilance
Earlsfort Terrace
IRL - Dublin 2
Tel: +353 1 6764971

Fax: +353 1 6762517
 Website: www.hpra.ie
 e-mail: medsafety@hpra.ie

4.9 Overdose

Symptoms of oral overdose may include nausea, vomiting, epigastric distress, diarrhoea and haematuria.

In the event of severe overdosage, general supportive care is recommended, including close clinical and laboratory monitoring of haematological, renal and hepatic functions, and coagulation status until the patient is stable. Forced diuresis, peritoneal dialysis, haemodialysis, or charcoal haemoperfusion have not been established as beneficial for an overdose of cefalexin. It would be extremely unlikely that one of these procedures would be indicated.

Unless 5 to 10 times the normal total daily dose has been ingested, gastro-intestinal decontamination should not be necessary.

There have been reports of haematuria without impairment of renal function in children accidentally ingesting more than 3.5g of cefalexin in a day. Treatment has been supportive (fluids) and no sequelae have been reported.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: First generation cephalosporins, ATC code: J01DB01

In vitro tests demonstrate that cephalosporins are bactericidal because of their inhibition of cell wall synthesis.

Cefalexin is active against the following organisms *in vitro*:

Beta-haemolytic streptococci

Staphylococci, including coagulase-positive, coagulase-negative and penicillinase-producing strains

Streptococcus pneumoniae

Escherichia coli

Proteus mirabilis

Klebsiella species

Haemophilus influenzae

Branhamella catarrhalis

Most strains of enterococci (*Streptococcus faecalis*) and a few strains of staphylococci are resistant to cefalexin. It is not active against most strains of *Enterobacter* species, *Morganella morganii* and *Pr. vulgaris*. It has no activity against *Pseudomonas* or *Herellea* species or *Acinetobacter calcoeticus*. Penicillin-resistant *Streptococcus pneumoniae* is usually cross resistant to beta lactam antibiotics. When tested by *in vitro* methods, staphylococci exhibit cross-resistance between cefalexin and methicillin-type antibiotics.

5.2 Pharmacokinetic properties

Cefalexin is acid stable.

Cefalexin is almost completely absorbed from the gastro-intestinal tract, and 75-100% is rapidly excreted in active form in the urine. Absorption is slightly reduced if the drug is administered with food. The half-life is approximately 60 minutes in patients with normal renal function. Haemodialysis and peritoneal dialysis will remove cefalexin from the blood.

Peak blood levels are achieved one hour after administration and therapeutic levels are maintained for 6-8 hours.

Approximately 80% of the active drug is excreted in the urine within 6 hours. No accumulation is seen with dosages above the therapeutic maximum of 4g/day.

The half-life may be increased in neonates due to their renal immaturity, but there is no accumulation when given at up to 50mg/kg/day.

5.3 Preclinical safety data

There are no pre-clinical data of relevance to the prescriber which are additional to that already included in other sections of the SPC.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

- Sucrose
- Imitation of Guarana Flavour
- Allura red AC
- Sodium laurilsulfate
- Methycellulose
- Dimeticone
- Xanthan gum
- Starch, pregelatinised

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

Unreconstituted product:	2 years
Bottle of reconstituted product:	10 days

6.4 Special precautions for storage

Powder: Do not store above 25°C.

Keep the bottle in the outer carton.
Keep the bottle tightly closed.

Suspension: After mixing the suspension should be stored at 2-8°C and used within 10 days.

6.5 Nature and contents of container

The powder is filled into 100 ml HDPE bottles with screw caps.

6.6 Special precautions for disposal and other handling

First invert the bottle and tap to loosen the powder then add a total of 60ml water in two portions, shaking after each addition until suspended. Where dilution is unavoidable, Syrup BP should be used after suspension has been prepared according to the manufacturers instructions.

The reconstituted product is coloured red.
Shake well before use.

7 MARKETING AUTHORISATION HOLDER

Flynn Pharma Limited
Alton House
4 Herbert Street
Dublin 2
Republic of Ireland

8 MARKETING AUTHORISATION NUMBER

PA 1226/2/2

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 14 April 1977

Date of last renewal: 14 April 2007

10 DATE OF REVISION OF THE TEXT

May 2017