

## Package leaflet: Information for the user

### Apixaban Clonmel 2.5 mg film-coated tablets

Apixaban

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### What is in this leaflet

1. What Apixaban Clonmel is and what it is used for
2. What you need to know before you take Apixaban Clonmel
3. How to take Apixaban Clonmel
4. Possible side effects
5. How to store Apixaban Clonmel
6. Contents of the pack and other information

#### 1. What Apixaban Clonmel is and what it is used for

Apixaban Clonmel contains the active substance apixaban and belongs to a group of medicines called anticoagulants. This medicine helps to prevent blood clots from forming by blocking Factor Xa, which is an important component of blood clotting.

Apixaban Clonmel is used in adults:

- to prevent blood clots (deep vein thrombosis [DVT]) from forming after hip or knee replacement operations. After an operation to the hip or knee, you may be at a higher risk of developing blood clots in your leg veins. This can cause the legs to swell, with or without pain. If a blood clot travels from your leg to your lungs, it can block blood flow causing breathlessness, with or without chest pain. This condition (pulmonary embolism) can be life threatening and requires immediate medical attention
- to prevent a blood clot from forming in the heart of patients with an irregular heartbeat (atrial fibrillation) and at least one additional risk factor. Blood clots may break off and travel to the brain and lead to a stroke, or to other organs and prevent normal blood flow to that organ (also known as a systemic embolism). A stroke can be life threatening and requires immediate medical attention
- to treat blood clots in the veins of your legs (deep vein thrombosis) and in the blood vessels of your lungs (pulmonary embolism), and to prevent blood clots from re-occurring in the blood vessels of your legs and/or lungs

#### 2. What you need to know before you take Apixaban Clonmel

**DO NOT take Apixaban Clonmel if you:**

- are **allergic** to apixaban or any of the other ingredients of this medicine (listed in section 6)
- are **bleeding excessively**
- have a **disease in an organ** of the body that increases the risk of serious bleeding (such as an **active or recent ulcer** of your stomach or bowel, **recent bleeding in your brain**)
- have a **liver disease** which leads to increased risk of bleeding (hepatic coagulopathy)
- are **taking medicines to prevent blood clotting** (e.g. warfarin, rivaroxaban, dabigatran or heparin), except when changing anticoagulant treatment; while having a venous or arterial line and you get heparin through this line to keep it open; or if a tube is inserted into your blood vessel (catheter ablation) to treat an irregular heartbeat (arrhythmia)

## Warnings and precautions

Talk to your doctor, pharmacist or nurse before you take this medicine if you have any of the following:

- an increased risk of bleeding, such as:
  - **bleeding disorders**, including conditions resulting in reduced platelet activity
  - **very high blood pressure**, not controlled by medical treatment
  - you are older than 75 years
  - you weigh 60 kg or less
- **severe kidney disease or if you are on dialysis**
- **a liver problem or a history of liver problems**
- this medicine will be used with caution in patients with signs of altered liver function
- **had a tube (catheter) or an injection into your spinal column** (for anaesthesia or pain reduction). Your doctor will tell you to take this medicine 5 hours or more after catheter removal
- **a prosthetic heart valve**
- your doctor determines that your blood pressure is unstable or another treatment or surgical procedure to remove the blood clot from your lungs is planned

Take special care with Apixaban Clonmel

- if you know that you have a disease called **antiphospholipid syndrome** (a disorder of the immune system that causes an increased risk of blood clots), tell your doctor who will decide if the treatment may need to be changed

If you need to have **surgery or a procedure which may cause bleeding**, your doctor might ask you to temporarily stop taking this medicine for a short while. If you are not sure whether a procedure may cause bleeding, ask your doctor.

## Children and adolescents

This medicine is not recommended in children and adolescents under 18 years of age.

## Other medicines and Apixaban Clonmel

Tell your doctor, pharmacist or nurse if you are taking, have recently taken or might take any other medicines.

Some medicines may increase the effects of Apixaban Clonmel and some may decrease its effects. Your doctor will decide if you should be treated with Apixaban Clonmel when taking these medicines and how closely you should be monitored.

The following medicines may increase the effects of Apixaban Clonmel and increase the chance for unwanted bleeding:

- some **medicines for fungal infections** (e.g. ketoconazole, etc.)
- some **antiviral medicines for HIV / AIDS** (e.g. ritonavir)
- other **medicines that are used to reduce blood clotting** (e.g. enoxaparin, etc.)
- **anti-inflammatory or pain medicines** (e.g. acetylsalicylic acid or naproxen). Especially, if you are older than 75 years and are taking acetylsalicylic acid you may have an increased chance of bleeding
- **medicines for high blood pressure or heart problems** (e.g. diltiazem)
- **antidepressant medicines called selective serotonin re-uptake inhibitors or serotonin norepinephrine re-uptake inhibitors**

The following medicines may reduce the ability of Apixaban Clonmel to help prevent blood clots from forming:

- **medicines to prevent epilepsy or seizures** (e.g. phenytoin, etc.)
- **St John's Wort** (a herbal supplement used for depression)
- **medicines to treat tuberculosis or other infections** (e.g. rifampicin)

## Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor, pharmacist or nurse for advice before taking this medicine.

### Pregnancy

The effects of apixaban on pregnancy and the unborn child are not known. You should not take this medicine if you are pregnant. **Contact your doctor immediately** if you become pregnant while taking this medicine.

### Breast-feeding

It is not known if apixaban passes into human breast milk. Ask your doctor, pharmacist or nurse for advice before taking this medicine while breast-feeding. They will advise you whether to stop breast-feeding or to stop/not start taking this medicine.

### **Driving and using machines**

Apixaban has not been shown to impair your ability to drive or use machines.

### **Apixaban Clonmel contains lactose and sodium**

**Lactose:** If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

**Sodium:** This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

### **3. How to take Apixaban Clonmel**

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor, pharmacist or nurse if you are not sure.

#### **Dose**

Swallow the tablet with a drink of water. Apixaban Clonmel can be taken with or without food. Try to take the tablets at the same times every day to have the best treatment effect.

If you have difficulty swallowing the tablet whole, talk to your doctor about other ways to take Apixaban Clonmel. The tablet may be crushed and mixed with water, or 5 % dextrose in water, or apple juice or apple puree, immediately before you take it.

#### **Instructions for crushing:**

- crush the tablets with a pestle and mortar
- transfer all the powder carefully into a suitable container then mix the powder with a little e.g., 30 ml (2 tablespoons), water or one of the other liquids mentioned above to make a mixture
- swallow the mixture
- rinse the pestle and mortar you used for crushing the tablet and the container, with a little water or one of the other liquids (e.g., 30 ml), and swallow the rinse

If necessary, your doctor may also give you the crushed Apixaban Clonmel tablet mixed in 60 ml of water or 5 % dextrose in water through a nasogastric tube.

#### **Take Apixaban Clonmel as recommended for the following:**

##### To prevent blood clots from forming after hip or knee replacement operations

The recommended dose is one tablet of Apixaban Clonmel **2.5 mg** twice a day. For example, one in the morning and one in the evening.

You should take the first tablet 12 to 24 hours after your operation.

If you have had a major **hip** operation, you will usually take the tablets for 32 to 38 days.

If you have had a major **knee** operation, you will usually take the tablets for 10 to 14 days.

##### To prevent a blood clot from forming in the heart in patients with an irregular heartbeat and at least one additional risk factor

The recommended dose is one tablet of Apixaban Clonmel **5 mg** twice a day.

The recommended dose is one tablet of Apixaban Clonmel 2.5 mg twice a day if:

- you have **severely reduced kidney function**
- **two or more of the following apply to you:**
  - your blood test results suggest poor kidney function (value of serum creatinine is 1.5 mg/dl (133 micromole/l) or greater)
  - you are 80 years old or older
  - your weight is 60 kg or lower

The recommended dose is one tablet twice a day, for example, one in the morning and one in the evening. Your doctor will decide how long you must continue treatment for.

#### To treat blood clots in the veins of your legs and blood clots in the blood vessels of your lungs

The recommended dose is **two tablets** of Apixaban Clonmel **5 mg** twice a day for the first 7 days, for example, two in the morning and two in the evening.

After 7 days, the recommended dose is **one tablet** of Apixaban Clonmel is **5 mg** twice a day, for example, one in the morning and one in the evening.

#### For preventing blood clots from re-occurring following completion of 6 months of treatment

The recommended dose is **one tablet** of Apixaban Clonmel **2.5 mg** twice a day for example, one in the morning and one in the evening.

Your doctor will decide how long you must continue treatment for.

#### **Your doctor might change your anticoagulant treatment as follows:**

- *Changing from Apixaban Clonmel to anticoagulant medicines*  
Stop taking Apixaban Clonmel. Start treatment with the anticoagulant medicines (for example heparin) at the time you would have taken the next tablet.
- *Changing from anticoagulant medicines to Apixaban Clonmel*  
Stop taking the anticoagulant medicines. Start treatment with Apixaban Clonmel at the time you would have had the next dose of anticoagulant medicine, then continue as normal.
- *Changing from treatment with anticoagulant containing vitamin K antagonist (e.g. warfarin) to Apixaban Clonmel*  
Stop taking the medicine containing a vitamin K antagonist. Your doctor needs to do blood measurements and instruct you when to start taking Apixaban Clonmel.
- *Changing from Apixaban Clonmel to anticoagulant treatment containing vitamin K antagonist (e.g. warfarin)*  
If your doctor tells you that you have to start taking the medicine containing a vitamin K antagonist, continue to take Apixaban Clonmel for at least 2 days after your first dose of the medicine containing a vitamin K antagonist. Your doctor needs to do blood measurements and instruct you when to stop taking Apixaban Clonmel.

#### **Patients undergoing cardioversion**

If your abnormal heartbeat needs to be restored to normal by a procedure called cardioversion, take this medicine at the times your doctor tells you, to prevent blood clots in blood vessels in your brain and other blood vessels in your body.

#### **If you take more Apixaban Clonmel than you should**

**Tell your doctor immediately** if you have taken more than the prescribed dose of this medicine. Take the medicine pack with you, even if there are no tablets left.

If you take more Apixaban Clonmel than recommended, you may have an increased risk of bleeding. If bleeding occurs, surgery, blood transfusions or other treatments that may reverse anti-factor Xa activity may be required.

### **If you forget to take Apixaban Clonmel**

- Take the dose as soon as you remember and:
  - take the next dose of Apixaban Clonmel at the usual time
  - then continue as normal.

**If you are not sure what to do or have missed more than one dose**, ask your doctor, pharmacist or nurse.

### **If you stop taking Apixaban Clonmel**

Do not stop taking this medicine without talking to your doctor first, because the risk of developing a blood clot could be higher if you stop treatment too early.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them. Apixaban Clonmel can be given for three different medical conditions. The known side effects and how frequently they occur for each of these medical conditions may differ and are listed separately below. For these conditions, the most common general side effect of this medicine is bleeding which may be potentially life threatening and requires immediate medical attention.

**The following side effects are known if you take apixaban to prevent blood clots from forming after hip or knee replacement operations.**

**Common** (may affect up to 1 in 10 people):

- anaemia which may cause tiredness or paleness
- bleeding including bruising and swelling
- nausea (feeling sick)

**Uncommon** (may affect up to 1 in 100 people):

- reduced number of platelets in your blood (which can affect clotting)
- bleeding:
  - occurring after an operation including bruising and swelling, blood or liquid leaking from the surgical wound/incision (wound secretion) or injection site
  - in your stomach, bowel or bright/red blood in the stools
  - blood in the urine
  - from your nose
  - from the vagina
- low blood pressure which may make you feel faint or have a quickened heartbeat
- blood tests may show:
  - abnormal liver function
  - an increase in some liver enzymes
  - an increase in bilirubin, a breakdown product of red blood cells, which can cause yellowing of the skin and eyes
- itching

**Rare** (may affect up to 1 in 1,000 people):

- allergic reactions (hypersensitivity) which may cause swelling of the face, lips, mouth, tongue and/or throat and difficulty breathing. **Contact your doctor immediately** if you experience any of these symptoms
- bleeding:
  - into a muscle
  - in your eyes
  - from your gums and blood in your spit when coughing
  - from your rectum

- hair loss

**Not known** (frequency cannot be estimated from the available data)

- bleeding:
  - in your brain or in your spinal column
  - in your lungs or your throat
  - in your mouth
  - into your abdomen or the space behind your abdominal cavity
  - from a haemorrhoid
  - tests showing blood in the stools or in the urine
- skin rash which may form blisters and looks like small targets (central dark spots surrounded by a paler area, with a dark ring around the edge) (*erythema multiforme*)
- blood vessel inflammation (vasculitis) which may result in skin rash or pointed, flat, red, round spots under the skin's surface or bruising.

**The following side effects are known if you take apixaban to prevent a blood clot from forming in the heart in patients with an irregular heartbeat and at least one additional risk factor.**

**Common** (may affect up to 1 in 10 people):

- bleeding:
  - in your eyes
  - in your stomach or bowel
  - from your rectum
  - blood in the urine
  - from your nose
  - from your gums
  - bruising and swelling
- anaemia which may cause tiredness or paleness
- low blood pressure which may make you feel faint or have a quickened heartbeat
- nausea (feeling sick)
- blood tests may show:
  - an increase in gamma-glutamyltransferase (GGT)

**Uncommon** (may affect up to 1 in 100 people):

- bleeding:
  - in your brain or spinal column
  - in your mouth or blood in your spit when coughing
  - into your abdomen, or from the vagina
  - bright/red blood in the stools
  - occurring after an operation including bruising and swelling, blood or liquid leaking from the surgical wound/incision (wound secretion) or injection site
  - from a haemorrhoid
  - tests showing blood in the stools or in the urine
- reduced number of platelets in your blood (which can affect clotting)
- blood tests may show:
  - abnormal liver function
  - an increase in some liver enzymes
  - an increase in bilirubin, a breakdown product of red blood cells, which can cause yellowing of the skin and eyes
- skin rash
- itching
- hair loss
- allergic reactions (hypersensitivity) which may cause swelling of the face, lips, mouth, tongue and/or throat and difficulty breathing. **Contact your doctor immediately** if you experience any of these symptoms

**Rare** (may affect up to 1 in 1,000 people):

- bleeding:
  - in your lungs or throat
  - into the space behind your abdominal cavity
  - into a muscle

**Very rare** (may affect up to 1 in 10,000 people):

- skin rash which may form blisters and looks like small targets (central dark spots surrounded by a paler area, with a dark ring around the edge) (*erythema multiforme*)

**Not known** (frequency cannot be estimated from the available data)

- blood vessel inflammation (vasculitis) which may result in skin rash or pointed, flat, red, round spots under the skin's surface or bruising.

**The following side effects are known if you take apixaban to treat or prevent re-occurrence of blood clots in the veins of your legs and blood clots in the blood vessels of your lungs.**

**Common** (may affect up to 1 in 10 people):

- bleeding:
  - from your nose
  - from your gums
  - blood in the urine
  - bruising and swelling
  - in your stomach or your bowel
  - from your rectum
  - in your mouth
  - from the vagina
- anaemia which may cause tiredness or paleness
- reduced number of platelets in your blood (which can affect clotting)
- nausea (feeling sick)
- skin rash
- blood tests may show:
  - an increase in gamma-glutamyltransferase (GGT) or alanine aminotransferase (ALT)

**Uncommon** (may affect up to 1 in 100 people):

- low blood pressure which may make you feel faint or have a quickened heartbeat
- bleeding:
  - in your eyes
  - in your mouth or blood in your spit when coughing
  - bright/red blood in the stools
  - tests showing blood in the stools or in the urine
  - occurring after an operation including bruising and swelling, blood or liquid leaking from the surgical wound/incision (wound secretion) or injection site
  - from a haemorrhoid
  - into a muscle
- itching
- hair loss
- allergic reactions (hypersensitivity) which may cause swelling of the face, lips, mouth, tongue and/or throat and difficulty breathing. Contact your doctor immediately if you experience any of these symptoms
- blood tests may show:
  - abnormal liver function
  - an increase in some liver enzymes
  - an increase in bilirubin, a breakdown product of red blood cells, which can cause yellowing of the skin and eyes

**Rare** (may affect up to 1 in 1,000 people):

- bleeding:
  - in your brain or in your spinal column
  - in your lungs

**Not known** (frequency cannot be estimated from the available data):

- bleeding:
  - into your abdomen or the space behind your abdominal cavity
- skin rash which may form blisters and looks like small targets (central dark spots surrounded by a paler area, with a dark ring around the edge) (*erythema multiforme*)
- blood vessel inflammation (vasculitis) which may result in skin rash or pointed, flat, red, round spots under the skin's surface or bruising.

### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist, or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via:

HPRA Pharmacovigilance

Website: [www.hpra.ie](http://www.hpra.ie)

By reporting side effects you can help provide more information on the safety of this medicine.

### **5. How to store Apixaban Clonmel**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and on the blister after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

### **6. Contents of the pack and other information**

#### **What Apixaban Clonmel contains**

The active substance is apixaban. Each tablet contains 2.5 mg of apixaban.

The other ingredients are:

- Tablet core: cellulose, microcrystalline, croscarmellose sodium, docusate sodium, polyvinyl caprolactam-polyvinyl acetate-polyethylene glycol graft copolymer, magnesium stearate.
- Film coat (opadry II yellow): hypromellose (2910, 15mPas), lactose monohydrate, titanium dioxide (E171), macrogol (3350), iron oxide yellow (E172).

#### **What Apixaban Clonmel looks like and contents of the pack**

The film-coated tablets are yellow, round and biconvex.

- They come in blisters in cartons of 10, 20, 30, 60, 100, 168 and 200 film-coated tablets.
- Unit dose blisters in cartons of 10 x 1, 20 x 1, 30 x 1, 60 x 1, 100 x 1, 168 x 1 and 200 x 1 film-coated tablets are also available.

Not all pack sizes may be marketed.

Patient Alert Card: handling information

Inside the Apixaban Clonmel pack together with the package leaflet you will find a Patient Alert Card or your doctor might give you a similar card.



This Patient Alert Card includes information that will be helpful to you and alert other doctors that you are taking Apixaban Clonmel. You should keep this card with you at all times.

1. Take the card
2. Separate your language as needed (this is facilitated by the perforated edges)
3. Complete the following sections or ask your doctor to do it:
  - Name:
  - Birth Date:
  - Indication:
  - Dose: mg twice daily
  - Doctor's name:
  - Doctor's telephone:
4. Fold the card and keep it with you at all times

#### **Marketing Authorisation Holder**

Clonmel Healthcare Ltd, Waterford Road, Clonmel, Co. Tipperary, Ireland

#### **Manufacturer**

STADA Arzneimittel AG, Stadastrasse 2-18, 61118 Bad Vilbel, Germany  
Clonmel Healthcare Ltd, Waterford Road, Clonmel, Co. Tipperary, Ireland

#### **This medicinal product is authorised in the Member States of the EEA under the following names:**

Germany:	Apixaban STADA 2,5 mg Filmtabletten
Austria:	Apixaban STADA 2,5 mg Filmtabletten
Belgium:	Apixaban EG 2,5 mg filmomhulde tabletten
Czech Republic:	Apixaban STADA
Denmark:	Apixaban STADA
Finland:	Apixaban STADA 2,5 mg kalvopäällysteiset tabletit
France:	APIXABAN EG 2,5 mg, comprimé pelliculé
Hungary:	Apixaban Stada 2,5 mg filmtabletta
Ireland:	Apixaban Clonmel 2.5 mg film-coated tablets
Iceland:	Apixaban STADA 2,5 mg filmuhúðuð tafla
Luxembourg:	Apixaban 2,5 mg and 5 mg EG comprimés pelliculés
The Netherlands:	Apixaban CF 2,5 mg, filmomhulde tabletten
Portugal:	Apixabano Ciclum
Romania:	Apixaban STADA 2,5 mg comprimate filmate
Slovenia:	Apiksaban STADA 2,5 mg filmsko obložene tablete
Slovakia:	Apixaban STADA 2,5 mg
Spain:	Apixiban STADA 2,5 mg comprimidos recubiertos con película EFG
Sweden:	Apixaban STADA 2,5 mg filmdragerade tabletter

**This leaflet was last revised in February 2022.**