



PACKAGE LEAFLET: INFORMATION FOR THE USER

85669.IG025
7.30095.0008/IRL

Ilvico® Cold and Flu film-coated tablets

Paracetamol 325mg, Caffeine 30mg, Brompheniramine maleate 3mg

Read all this leaflet carefully because it contains important information for you.

This medicine is available without prescription. However, you still need to use Ilvico carefully to get the best results from it. Ilvico Cold and Flu film-coated tablets will be referred to as Ilvico throughout the leaflet

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- You must contact a doctor if your symptoms worsen or do not improve.
- If any of the side effects get serious, or if you notice any side effect not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

1. What Ilvico is and what it is used for
2. Before you use Ilvico
3. How to use Ilvico
4. Possible side effects
5. How to store Ilvico
6. Further information

1. WHAT ILVICO IS AND WHAT IT IS USED FOR

Ilvico Cold and Flu film-coated tablets contain paracetamol, caffeine and brompheniramine. Ilvico Cold and Flu film-coated tablets are used for the relief of symptoms associated with the common cold, influenza and upper respiratory tract infections.

2. BEFORE YOU TAKE ILVICO TABLETS

Do not take Ilvico

- if you are allergic (hypersensitive) to paracetamol, caffeine or brompheniramine, or any of the other ingredients of Ilvico.
- You suffer from severe liver diseases (with or without liver insufficiency), viral hepatitis (increases the risk of liver toxicity)
- You suffer from Peptic ulcers
- Your suffer from Severe kidney impairment
- You suffer from Haemophilia (a disorder that affects blood clotting)
- You are less then 12 years old
- If you are pregnant or are breast-feeding
- If you have tachyarrhythmia (fast heart rate)
- If you suffer from narrow-angle glaucoma
- If you suffer from brain damage or epilepsy

Take special care with Ilvico if

- You suffer from anaemia or you have heart, lung, liver, or kidney problems. Do not take Ilvico for long periods of time.
- You drink more than three alcoholic drinks a day. Ilvico may cause liver damage.
- You are an alcoholic. You need to be careful how much paracetamol you take - talk to your doctor or pharmacist.
- You are undergoing diagnostic tests involving blood and/or urine, as Ilvico may interfere with these tests.
- As brompheniramine may interfere with allergenic tests on the skin, you should suspend treatment with Ilvico at least 3 days before taking such tests.
- You have aspirin sensitive asthma. This may get worse when taking these tablets.
- You have a bladder problem called vesical neck obstruction. This may get worse when taking these tablets.
- You have an enlarged prostate or urinary retention. Your symptoms may get worse when taking these tablets.

Taking other medicines

This product CONTAINS PARACETAMOL and DO NOT TAKE ANY OTHER PARACETAMOL CONTAINING PRODUCTS.

Please tell your doctor or phamacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription, specifically the following:

- Anticoagulants (to prevent blood clotting e.g. warfarin or acenocoumarol)
- Medicines containing alcohol (including some cold remedies)
- Anti-convulsants (to prevent seizures and epilepsy e.g. phenytoin, phenobarbital, methyl phenobarbital, primidone, lamotrigine)
- Medicines containing oestrogen such as oral contraceptives ('the pill') and hormone replacement therapy (HRT)
- Antibiotics such as rifampicin, chloramphenicol and isoniazid
- Anticholinergic drugs such as glycopyrrolate and propanteline
- Antiretroviral drugs for treatment of HIV infection such as zidovudine (ATZ)
- Antidepressants such as Monoamine Oxidase Inhibitors (MAOIs) e.g. phenelzine and isocarboxazid
- Other antidepressants such as tricyclic anti-depressants e.g. imipramine, lofepramine, doxepin
- Barbiturates e.g. phenobarbital
- Other paracetamol containing medicines, or other analgesic drugs (painkillers)
- Antimuscarinic drugs such as propantheline bromide used to treat cramps or spasms of the stomach or excessive sweating
- Antiemetic (prevention of feeling sick) drugs such as metoclopramide

Always tell your doctor or pharmacist about all the medicines you are taking. This means medicines you have bought yourself as well as those you have on prescription from your doctor, such as probenecid (a medicine for gout), propranolol (a medicine for your heart or treatment of blood pressure), loop diuretics (water tablets), cholestyramine (a medicine for high cholesterol levels or itching caused by excess bile) or drugs prescribed for insomnia (difficulty sleeping).

Taking Ilvico with food and drink

Avoid drinking alcohol when taking Ilvico Tablets. The tablets should be swallowed with a drink of water preferably after a meal.

Pregnancy and breast-feeding

Do not take Ilvico tablets when pregnant or breast-feeding unless advised by a doctor.

Driving and using machines

Ilvico Tablets may make you feel drowsy. If you are affected you should not drive or operate machinery.

Important Information about some of the ingredients of Ilvico

Suitable for people with coeliac disease.

3. HOW TO TAKE ILVICO

The product should only be used when clearly necessary. You should check with your doctor or pharmacist if you are not sure. Do not exceed the recommended dose.

You should not take Ilvico and Flu Film-Coated Tablets with other paracetamol-containing products.

Continued, please turn over.





IMMEDIATE MEDICAL ADVICE SHOULD BE SOUGHT IN THE EVENT OF OVERDOSAGE EVEN IF YOU FEEL WELL BECAUSE OF THE RISK OF IRREVERSIBLE LIVER DAMAGE.

The usual dose is:

Adults and children older than 12 years

One or two tablets taken orally with water three times daily.
Long term use without medical supervision may be harmful.

You should stop taking this medicine once your symptoms have disappeared unless you have any problems. In that case, check with your doctor. If symptoms persist or worsen you should consult your doctor.

If you take more Ilvico that you should

Contact your doctor or go to your nearest hospital casualty department immediately, even if you feel well, because of the risk of delayed, serious liver damage.
Take your tablets or the pack with you so that the doctor knows what you have taken.

If you forget to take Ilvico

If you have missed a dose, take it as soon as you remember and then take your next dose after 8 hours. Do not take a double dose to make up for a forgotten dose.

4. POSSIBLE SIDE EFFECTS

Like all medicines, Ilvico Cold and Flu Film-Coated Tablets can cause side effects, although not everybody gets them.
If any of the side effects get serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

This product contains an antihistamine, which has been known to cause, in children and occasionally in adults, insomnia (difficulty in sleeping), nervousness, high fever, tremors and fits.

Allergic reactions which can include skin rash, itching, bruising, painful red areas, peeling or blistering, dizziness, difficulty in breathing, wheezing or shortness of breath, swollen, face throat, lips, hands or fingers - these have been reported very rarely (less than 1 in 10,000 of users).

The following side effects have been reported frequently (between 1 in 10 and 1 in 100 users):

- Drowsiness
- Other undesirable effects include dizziness, lack of muscular coordination, numbness and tingling, headache and restlessness.
- A dry mouth
- High blood pressure (hypertension)

The following side effects have been reported rarely (between 1 in 1,000 and 1 in 10,000 of users):

- Feeling sick (nausea)
- Excitation
- Low blood pressure (hypotension)

The following side effects have been reported very rarely (less than 1 in 10,000 of users):

- Blood disorders such as low red and/or white blood cells and/or platelet levels.
Symptoms include tiredness, fatigue, bruising of skin, nose bleeds, bleeding gums, infections, fever, throat pain, ulcers in the throat and mouth.
- Bronchospasm (difficulty in breathing) or dyspnoea (shortness of breath)
- Insomnia (unable to sleep)
- Nervousness, hyperpyrexia (high fever)
- Tremors or convulsions
- Skin disorders such as blisters or non specific skin reactions and sweating.

If you notice any unwanted effects, even ones not mentioned here, please go and see your doctor.

5. HOW TO STORE ILVICO

Keep out of the sight and reach of children.
Do not store above 25°C. Do not use the tablets after the expiry date shown on the carton and blister strip.
Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. FURTHER INFORMATION

What Ilvico contains

- The active substance(s) are paracetamol 325mg, caffeine 30mg and brompheniramine maleate 3mg.
- The other ingredient(s) are microcrystalline cellulose, sodium carboxymethylstarch Type A, crospovidone, hypromellose, cellulose powder, magnesium stearate, titanium dioxide E171, glycerol, macrogol 4000, macrogol 6000, talc, colloidal anhydrous silica.

What Ilvico looks like and contents of the pack

The tablets are almost white, round film-coated tablets. They are supplied in blister packs containing 20 tablets.

Marketing Authorisation Holder

Seven Seas Limited, Bedfont Cross, Stanwell Road, Feltham, TW14 8NX, United Kingdom.

Manufacturer

Merck Farma y Quimica, S.L., Poligono MERCK, 08100 Mollet del Valles, Barcelona, Spain, or
Merck KGaA & Co., Werk Spittal, Hösslgasse 20, 9800 Spittal/Drau, Austria.

Product Authorisation Number:
PA : 417/18/1

Date of revision: July 2015

If you would like the leaflet in a different format, please contact the license holder at the above address.

Ilvico Irland

Leaflet 20

Format (A x B):	175 x 297 mm	Reference-No. MSM:	7.30095.0008
Fontsize:	8,5 pt	Article-No. MSM:	30095
Program:	Illustrator CS4	Manufacturer-No.:	85669.G025
Manufacturer:	Merck Spittal	Laetuscode:	13478-2121121211222
Way of dispatch:	To Merck Spittal, Mrs. Aichholzer		
Printing colours:	Black		
	-		
Non printing colours:	Cutter		
Release msm:	Approved 23.07.15 Mrs. Milligan, Mr. Morton		
Technical release:	Approved 16.07.15 Mrs. Luft		
Designed by: K_150104	Jürgen Kiffe • KIFFE GRAFIK DESIGN • In der Strieth 7 • 64395 Brensbach Fon 06161-5879728 • Mobil 0174-4129801 • grafik@kiffe-design.de		

Merck logo version: small	width: - mm	distance: double-M
Release Merck logo version:	-	